



## Vestibular Testing Information

Your physician has ordered a Videonystagmography (VNG) test to evaluate your balance system. First, you will have a diagnostic hearing evaluation. Then, the balance system of the ear and the brain will be evaluated by recording eye movement in response to changes in head position, viewing moving objects, motion of the body, and thermal stimuli (warm and cold) applied to the ear canal. Special cameras are used to measure these movements. A more detailed explanation will be given during each portion of the test. The testing will take approximately 1.5 to 2 hours.

**PLEASE NOTE: Cancellations must be made at least 48 hours prior to your appointment or a fee of \$25.00 will be charged to you directly.**

Your test has been scheduled for \_\_\_\_\_

Your test will be performed by \_\_\_\_\_

Please report to:

At this time \_\_\_\_\_

(Brighton- Clinton Crossings)

(Greece - Location)

CLEAR CHOICE HEARING AND BALANCE  
1065 SENATOR KEATING BLVD.  
SUITE 210  
ROCHESTER, NY 14618  
(585) 342-4327

CLEAR CHOICE HEARING AND BALANCE  
103 CANAL LANDING BLVD.  
SUITE 3  
ROCHESTER, NY 14626  
(585) 723-3440

Please follow the instructions carefully in order to insure accuracy of the test results and aid in your comfort during the testing. Please consult your physician regarding any questions you may have.

### 1. Medications

\*\*\*Please do not take the following medications more than 48 prior to the test time:

**Anti-dizziness Medications**

Dramamine, Bonine, Phenergan, Compazine, Antivert, Meclizine, Transderm, Scopalamine

**Tranquilizers**

Valium, Diazepam, Librium, Elavil, Triavil, Klonopin, Clonazepam, Xanax, Ativan

**Any Sleeping Pills**

Seconal, Dalmane, Nembutal, Phenobarbital\*, Tuinal

**Anti-depressants**

Nortriptyline, Paxil, Prozac, St. John's Wort, Celexa, Zoloft

**Pain Medication**

Codeine, Demerol, Percocet and Percodan

**Antihistamines**

Dimatane, Benadryl, Actifed, Dimetapp, Zyrtec, Allegra, Claritin or and cold/allergy pills

\*Please note that **phenobarbital should not be taken for 7-14 days** before the test.

\*\*\*Please note that any medications that are necessary to your long term and immediate health should continue to be taken. Examples include any heart, blood pressure, blood thinning, or epilepsy medications. **If you have any questions regarding medications, check with your physician.**

2. Please **do not eat or drink** anything for **3 hours** before your appointment.

\*\*Please **do not eat breakfast** or eat anything other than a drink of water if you are scheduled for a **morning appointment**. If your **appointment is in the afternoon**, please have a light breakfast and **do not have lunch**.

3. No alcohol for **48 hours** before the test.

4. No coffee, tea, or cola drinks the day of the exam.

5. Please **do not wear eye/facial makeup or lotions/creams** the day of the test.

6. Please **wear comfortable slacks/pants** the day of the test.

7. Please **do not wear contact lenses** the day of the test. Bring your glasses.

8. Please note that the table used for vestibular testing **supports body weight up to 400 lbs**. Please contact our office if this is a problem or concern.

9. While you probably will not experience any ill effects from the test, **you must have someone accompany you who can drive you home.**

\*\*\*Please fill out the attached **Dizziness and Balance Questionnaire** and bring it with you on the day of your test.

Greece

103 Canal Landing Blvd Ste. 3

Rochester, NY 14626

P: (585)723-3440

F: (585)735-4632

Brighton

1065 Senator Keating Blvd Ste 210

Rochester, NY 14618

P: (585)342-4327

F: (585)735-4632